

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 31/47, 9/107	A1	(11) International Publication Number: WO 99/61025 (43) International Publication Date: 2 December 1999 (02.12.99)
(21) International Application Number: PCT/EP99/03623 (22) International Filing Date: 26 May 1999 (26.05.99) (30) Priority Data: 9811200.6 26 May 1998 (26.05.98) GB 9818105.0 19 August 1998 (19.08.98) GB (71) Applicant (for all designated States except AT US): NOVARTIS AG [CH/CH]; Schwarzwaldallee 215, CH-4058 Basel (CH). (71) Applicant (for AT only): NOVARTIS-ERFINDUNGEN VERWALTUNGSGESELLSCHAFT MBH [AT/AT]; Brunner Strasse 59, A-1230 Vienna (AT). (72) Inventors; and (75) Inventors/Applicants (for US only): LANG, Steffen [DE/CH]; Predigerweg 1, CH-4153 Reinach (CH). LIECHTI, Kurt [CH/CH]; Bienenstrasse 6, CH-4104 Oberwil (CH). (74) Agent: BECKER, Konrad; Novartis AG, Corporate Intellectual Property, Patent & Trademark Dept., CH-4002 Basel (CH).		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: MICROEMULSION PRECONCENTRATES CONTAINING A PIPERIDINE SUBSTANCE P ANTAGONIST (57) Abstract Microemulsion comprising a piperidine, e.g. 1-acyl-piperidine Substance P Antagonist and its use in the treatment of CNS disorders, e.g. depression and social phobia, and respiratory diseases, e.g. asthma and chronic bronchitis.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

MICROEMULSION PRECONCENTRATES CONTAINING A PIPERIDINE SUBSTANCE P ANTAGONIST

The present invention relates to novel pharmaceutical compositions in which the active agent is a piperidine substance P antagonist, in particular a N-benzoyl-2-benzyl-4-(azanaphthoyl-amino) piperidine, useful for treatment and prevention of e.g. central nervous system disorders, e.g. depression, social phobia, or respiratory diseases, e.g. asthma and chronic bronchitis. 1-Acylpiperidine substance P antagonists are a class of compounds described e.g. in published European patent EP 0532456B1, the contents of which publication is incorporated herein by reference. Similarly, N-benzoyl-2-benzyl-4-(azanaphthoyl-amino) piperidines and their activity as Substance P Antagonists are described in published European patent application EP 0707006A, the contents of which application is incorporated herein by reference.

Piperidine substance P antagonists, such as disclosed in EP 0532456B1 and EP 0707006A, present highly specific difficulties in relation to administration generally and galenic compositions in particular, including in particular problems of drug bioavailability and variability in inter- and intra-patient dose response, necessitating development of a non-conventional dosage form.

In accordance with the present invention it has now surprisingly been found that stable pharmaceutical compositions with Piperidine substance P antagonists, having particularly interesting bioavailability characteristics and reduced variability in inter- and intra-subject bioavailability parameters, are obtainable. These novel compositions have been found to meet or substantially reduce the difficulties encountered previously. It has been shown that the compositions in accordance with the present invention may enable effective dosaging with concomitant enhancement as well as reduced variability of resorption/bioavailability levels for and between individual patients. Thus, the invention may achieve effective therapy with tolerable dosage levels of such Piperidine substance P antagonists, and may permit closer standardization and optimization of daily dosage requirements for each individual. Consequently, occurrence of potential undesirable side-effects is diminished and overall cost of

therapy may be reduced.

In one aspect the present invention provides a spontaneously dispersible pharmaceutical composition comprising a Piperidine substance P antagonist, hereinafter also referred to as the
5 active agent.

Such spontaneously dispersible pharmaceutical compositions are preferably in liquid form , or are produced from at least one liquid component and preferably are micellar precursors.

10 The "term spontaneously dispersible pharmaceutical composition" as used herein is defined as a system that is capable of producing colloidal structures, e.g. solid or preferably liquid particles, e.g. droplets and/or nanoparticles, and/or micellar compositions, e.g. aqueous micelles, e.g. of up to 1000 nm in diameter, when diluted with an aqueous medium, for example, in water, for example on dilution at a dilution ratio e.g. 1:10 of composition to water, or in gastric juices,
15 e.g. simulated conditions after oral application at a similar dilution ratio at e.g from 15 to 37 degrees Centigrade. The spontaneously dispersible pharmaceutical compositions are hereinafter also referred to as compositions of the invention. The colloidal structures may form spontaneously or substantially spontaneously when the components of the composition of the invention are brought into contact with an aqueous medium, e.g. by simple shaking by hand for
20 a short period of time, for example for 10 seconds. The compositions of the invention are thermodynamically stable, e.g. for at least 15 minutes or up to 4 hours, even to 24 hours. Typically, they contain dispersed structures of a mean diameter less than about 300 nm (3,000 Å), e.g. less than about 200 nm (2,000 Å) as measured by standard light scattering techniques, e.g. using a Malvern Zetasizer 3000. Naturally the particles of mean diameter greater than 200
25 nm may be present, but these are preferably less than 50 % by weight, e.g. less than 10-20 % per weight, of the total. Preferably they comprise droplets or nanoparticles having a mean diameter of less than about 150 nm (1,500 Å), typically less than 100 nm (1,000 Å), generally greater than 5 nm (50 Å). Alternatively, the spontaneously dispersible pharmaceutical compositions may form upon dilution simultaneously a mixture comprising micelles and

nanoparticles. The micelles may be essentially monophasic and substantially non-opaque, i.e. transparent or opalescent when viewed by optical microscopic means. It was found that the proportion of nanoparticles present may be temperature dependent but still adequate bioavailability characteristics may be obtained.

5

In another aspect the present invention provides a spontaneously dispersible pharmaceutical composition comprising a piperidine substance P antagonist as the active agent, and a carrier medium comprising

- 10 1) a hydrophilic component, and
2) a surfactant.

Preferably such spontaneously dispersible pharmaceutical compositions are for oral administration. Conveniently the Piperidine substance P antagonist, is water insoluble, e.g. has
15 a solubility of below 0.001%, e.g. 0.001 to 0.0001 %. Conveniently the active agents are used in free base form.

The active agents may show Substance P antagonistic activity as indicated in standard in vitro or in vivo tests e.g. as disclosed in the above-mentioned European filings.

- 20 The piperidine may be a piperidine derivative which is an analogue or may be poly (e-g. di or tri-) substituted. The active agents include not only those disclosed in the above mentioned European filings but also analogues for example
- a) 1-[2-[3-(3,4-Dichlorophenyl)-1-[(3-(1-isopropoxyphenyl)acetyl)piperidin-3-yl]ethyl]-4-phenyl-1-azoniabicyclo[2.2.2]octane,
 - 25 b) (2S-cis)- N-[(2-methoxyphenyl)methyl]-2-phenyl-)piperidinamine [CP-99994],
 - c) cis-3-((3-methyl-5-trimethylsilyl)benzyloxy)-2-phenyl piperidine,
 - d) 4-phenyl-4-((1-(3,5-bis(trifluoromethyl)phenyl)-2-hydroxyethoxy)methyl)piperidine [L-733060],

- e) 2-phenyl-3-(3,5-bis(trifluoromethyl)benzyloxy)piperidine [L-733060],
 - f) (R)-3-(1H-indol-3-yl)-1-[N-(2-methoxybenzyl)acetylamino]-2-[N-(2-(4-(piperidin-1-yl)piperidin-1-yl)acetyl)amino]propane,
 - g) (R)-3-(1H-indol-3-yl)-1-[N-(2-methoxybenzyl)acetylamino]-2-[N-(2-(4-(piperidin-1-yl)piperazin-1-yl)acetyl)amino]propane,
 - h) (2-Methoxy-5-(5-trifluoromethyl-tetrazol-1-yl)-benzyl)-(2S-phenyl-1-piperidin-3S-yl)-amine [GR-205171],
 - i) (-)-(B)-cis-4-[1-[3,5-bis(trifluoromethyl)benzoyl]-2-(phenylmethyl)-4-piperidinyl]-N-(2,6-dimethylphenyl)-1-piperazine acetamide,
- 10 including salts thereof.

The active agents are preferably 1-acylpiperidine substance P antagonists e.g. as claimed in EP 0532456 B. Preferred compounds of the invention are the compounds of EP 0707006 A, e.g. N-benzoyl-2-benzyl-4-(azanaphthoyl-amino) piperidines, and most preferably (2R,4S)-N-(1-
 15 (3,5-bis(trifluoromethyl)-benzoyl)-2-(4-chlorobenzyl)-4-piperidinyl)-quinoline-4-carboxamide, hereinafter referred to as Compound A, as disclosed in EP 707 006.

The composition of the invention may also comprise a carrier medium containing further components. Typical further components of the compositions of the invention are, e.g.
 20 described in Fiedler, H. P. "Lexikon der Hilfsstoffe für Pharmazie, Kosmetik und angrenzende Gebiete", Editio Cantor, D-7960 Aulendorf, 4th revised and expanded edition (1996).

In accordance with the present invention the active agent may be present in an amount by weight of up to about 20% by weight of the composition of the invention, e.g. from about
 25 0.05% by weight. The active agent are preferably present in an amount of 0.5 to 15 % by weight of the composition, for example about 1 to 10 %, preferably 1.5 to 2% .

The hydrophilic component typically has a solubility in water of at least 1 g/100 ml or more, e.g. at least 5 g/100 ml at 25 degrees centigrade. It preferably provides for fast mixing of an

active agent with water. Such mixing may be determined by routine experimentation, for example by various chromatography methods, e.g. Gas Chromatography (GC). Conveniently the hydrophilic component may also be miscible with an organic solvent, e.g. ether. Preferably the hydrophilic component may be an alcohol, e.g. a water miscible alcohol, e.g. absolute
5 ethanol, or glycerol. Other alcohols include glycols, e.g. any glycol obtainable from an oxide such as ethylene oxide, e.g. 1,2-propylene glycol. Other examples are polyols, e.g. a polyalkylene glycol, e.g. poly(C₂₋₃)alkylene glycol. A typical example is a polyethylene glycol, e.g. of a preferred molecular weight of 200-1000 daltons, more preferably, 200-400 daltons, especially 400. Alternatively the hydrophilic component may preferably comprise an N-
10 alkylpyrrolidone, e.g. N-(C₁₋₁₄alkyl)pyrrolidone, e.g. N-methylpyrrolidone, tri(C₁₋₄alkyl)citrate, e.g. triethylcitrate, dimethylisosorbide, (C₅-C₁₃)alkanoic acid, e.g. caprylic acid or propylene carbonate. Most preferred is triethylcitrate or propylene glycol.

The hydrophilic component may comprise a main or sole component, e.g. an alcohol, e.g. C₁₋₄-
15 alcohol, e.g. ethanol, or alternatively a co-component, e.g. which may be selected from partial lower ethers or lower alkanols. Especially preferred partial ethers are those known and commercially available, for example Transcutol (which has the formula C₂H₅-[O-(CH₂)₂]-OH), Glycofurol (also known as tetrahydrofurfuryl alcohol polyethylene glycol ether), or lower alkanols such as ethanol. Preferred alkanol components include ethanol, 1,2-propylene glycol
20 or a polyethylene glycol, e.g. of a molecular weight of 100 to 600 daltons, e.g. polyethylene glycol 400. When present ethanol may comprise 0, e.g. 1 to, to 60 % by weight of the hydrophilic component; preferably 20 to about 55% by weight and more preferably about 40 to about 50 % by weight. The hydrophilic component may conveniently contain ethanol as sole component. The ratio of hydrophilic component to hydrophilic co-component is typically from
25 about 0.5:1 to about 2:1.

The hydrophilic component may comprise 5 to 50 % by weight of the composition of the invention, e.g. 10 to 50%; preferably 15 to 40 % by weight, more preferably about 20 to about 30 % by weight, e.g. from 15 to 35 % by weight.

Examples of suitable surfactants, e.g. preferably surfactants of high HLB value, e.g. HLB>10, for use in this invention are:

- 5 (i) Reaction products of a natural or hydrogenated castor oil and ethylene oxide, i.e. polyoxyethylene glycolated natural or hydrogenated vegetable oils, for example polyoxyethylene glycolated natural or hydrogenated castor oils. The natural or hydrogenated castor oil may be reacted with ethylene oxide in a molar ratio of from about 1:35 to about 1:60, with optional removal of the polyethyleneglycol component from the
- 10 products. Various such surfactants are commercially available. The polyethyleneglycol-hydrogenated castor oils available under the trade name CREMOPHOR are especially suitable. Particularly suitable are CREMOPHOR RH 40, which has a saponification number of about 50 to 60, an acid number less than about 1, a water content (Fischer) less than about 2%, an n_D^{60} of about 1.453 to 1.457 and an HLB
- 15 of about 14 to 16; and CREMOPHOR RH 60, which has a saponification number of about 40 to 50, an acid number less than about 1, an iodine number of less than about 1, a water content (Fischer) of about 4.5 to 5.5%, an n_D^{25} of about 1.453 to 1.457 and an HLB of about 15 to 17. An especially preferred product of this class is CREMOPHOR RH40.
- Also suitable are polyethyleneglycol castor oils such as that available under the trade name
- 20 CREMOPHOR EL, which has a molecular weight (by steam osmometry) of about 1630, a saponification number of about 65 to 70, an acid number of about 2, an iodine number of about 28 to 32 and an n_D^{25} of about 1.471 and an HLB value of about 12 to 14. Similar or identical suitable products which may also be used are the various tensides available under the trade names NIKKOL (e.g. NIKKOL HCO-40 and HCO-60), MAPEG (e.g. MAPEG
- 25 CO-40h), INCROCAS (e.g. INCROCAS 40), and TAGAT (for example polyoxyethylene-glycerol-fatty acid esters e.g. TAGAT RH 40. A preferred polyethoxylated glyceride is TAGAT TO, a polyoxyethylene-glycerol-trioleate having a HLB value of 11.3. These surfactants are further described in Fiedler loc. cit., incorporated herein by reference.

- (ii) Related products that belong to the class of polyoxyethylene alkyl ethers are available under the tradename BRIJ, e.g. Brij 35 which has an HLB value of about 16.9.
- 5 (iii) Polyoxyethylene-sorbitan-fatty acid esters (also called polysorbates), e.g. of from 4 to 25 alkylene moieties, for example mono- and tri-lauryl, palmityl, stearyl and oleyl esters of the type known and commercially available under the trade name TWEEN (Fiedler, loc.cit. p.1300-1304, incorporated herein by reference) including the products TWEEN
- 20 [polyoxyethylene(20)sorbitanmonolaurate],
- 10 21 [polyoxyethylene(4)sorbitanmonolaurate],
- 40 [polyoxyethylene(20)sorbitanmonopalmitate],
- 60 [polyoxyethylene(20)sorbitanmonostearate],
- 65 [polyoxyethylene(20)sorbitantristearate],
- 80 [polyoxyethylene(20)sorbitanmonooleate],
- 15 81 [polyoxyethylene(5)sorbitanmonooleate],
- 85 [polyoxyethylene(20)sorbitantrioleate].

Especially preferred products of this class are TWEEN 40 and TWEEN 80.

- 20 (iv) Polyoxyethylene fatty acid esters, for example polyoxyethylene stearic acid esters of the type known and commercially available under the trade name MYRJ (Fiedler, loc. cit., 2, p.834-835, incorporated herein by reference). An especially preferred product of this class is MYRJ 52 having a n_D^{50} of about 1.1, a melting point of about 40 to 44°C, an HLB value of about 16.9, an acid value of about 0 to 1 and a saponification no. of about 25 to
- 25 35. Other related products include polyethoxylated saturated hydroxy fatty acids which may be produced by reacting a saturated hydroxy fatty acid, e.g. C_{18} to C_{20} with e.g. ethylene oxide or polyethylene glycol. Suitable examples for the present invention include are known and commercially available, e.g. from the BASF company under the trade mark Solutol. Especially preferred is Solutol HS15 which is known, e.g. from the BASF

technical leaflet MEF 151E (1986), to comprise of about 70% polyethoxylated 12-hydroxystearate by weight and about 30% by weight unesterified polyethylene glycol component. Solutol HS 15 has a hydrogenation value of 90 to 110, a saponification value of 53 to 63, an acid number of maximum 1, and a maximum water content of 0.5% by weight.

- 5
- (v) Polyoxyethylene-polyoxypropylene co-polymers and block co-polymers, for example of the type known and commercially available under the trade names PLURONIC, EMKALYX and POLOXAMER (Fiedler, loc. cit., 2, p. 959, incorporated herein by reference). An especially preferred product of this class is PLURONIC F68, having a melting point of about 52°C and a molecular weight of about 6800 to 8975. A further preferred product of this class is POLOXAMER 188, which has an HLB value of about 29.
- 10
- (vi) Propylene glycol mono- and di-fatty acid esters such as propylene glycol dicaprylate (also known and commercially available under the trade name MIGLYOL 840), propylene glycol dilaurate, propylene glycol hydroxystearate, propylene glycol isostearate, propylene glycol laurate, propylene glycol ricinoleate, propylene glycol stearate and so forth (Fiedler, loc. cit., 2, p. 808-809, incorporated herein by reference). Propylene glycol mono C₈ esters include Sefsol 218 (Nikko Chemicals) and Capryol 90 (Gattefossé).
- 15
- (vii) Tocopherol esters, e.g. tocopheryl acetate and tocopheryl acid succinate (HLB of about 16).
- 20
- Further ionic surfactant classes not represented by the categories described above include
- (i) Docusate salts, for example dioctylsulfosuccinate or related compounds, for example di-[2-ethylhexyl]-succinate (Fiedler, loc. cit., 1, p. 107-108, incorporated herein by reference).

(ii) Phospholipids, in particular lecithins (Fiedler, loc. cit., 2, p. 943-944, incorporated herein by reference). Suitable lecithins include, in particular, soya bean lecithins.

5 The surfactant selected preferably has a hydrophilic-lipophilic balance (HLB) of at least 10, for example Cremophor, e.g. Cremophor RH40. A group of surfactants comprises those with a polyoxyethylene moiety.

The surfactant may comprise 5 to 80 % by weight of the composition of the invention;
10 preferably 10 to 70 % by weight, more preferably 20 to 60 % by weight and even more preferably between about 40 % and 55% by weight.

The compositions of the invention may further comprise a lipophilic phase or lipophilic component. These compositions may be capable of producing emulsions as colloidal
15 structures, preferably an aqueous microemulsion as a micelle, upon mixing with an aqueous medium.

In another aspect the present invention provides a spontaneously dispersible pharmaceutical composition, preferably in the form of a microemulsion preconcentrate, comprising a piperidine
20 substance P antagonist.

In yet another aspect the present invention provides a spontaneously dispersible pharmaceutical composition, preferably in the form of a microemulsion preconcentrate, comprising a piperidine substance P antagonist, e.g. Compound A, and a carrier medium comprising

25

- 1) a hydrophilic phase,
- 2) a lipophilic phase, and
- 3) a surfactant.

Such compositions of the invention may be in the form of "microemulsion preconcentrates" of the type providing o/w (oil-in-water) microemulsions. A "microemulsion preconcentrate" is defined in this specification as being a composition which spontaneously forms a microemulsion
5 in an aqueous medium, for example, in water, for example on dilution of 1:1 to 1:10, e.g. 1:10 or in the gastric juices after oral application. Typically, a "microemulsion" is a slightly opaque, opalescent, non-opaque or substantially non-opaque colloidal dispersion that is formed spontaneously or substantially spontaneously when its components are brought into contact with an aqueous medium. A microemulsion is thermodynamically stable and contains dispersed
10 particles of a mean diameter less than about 2000 Å. Generally microemulsions comprise droplets or liquid nanoparticles having a mean diameter of less than about 1500 Å; typically less than 100 nm, generally greater than 10 nm, and stable over periods in excess of 24 hours.

Naturally, such compositions of the invention comprising a hydrophilic phase, lipophilic phase
15 and surfactant may, e.g. before dilution, e.g. be preferably a hydrophilic component, lipophilic component and surfactant in the form of essentially a single phase system, e.g. a w/o microemulsion or other micellar composition.

Such compositions of the invention additionally containing an aqueous phase, preferably water,
20 may be in the form of a microemulsion.

The lipophilic component, when present, may preferably be characterized by a low HLB value of less than 10, e.g. up to 8, and may comprise (i) medium chain fatty acid triglycerides, (ii) mixed mono-, di-, tri-glycerides, (iii) transesterified ethoxylated vegetable oils, (iv) mixtures of
25 mono- and di-glycerides, or pure or substantially pure mono- or di-glycerides, (v) sorbitan fatty acid esters, (vi) pentaerythriol fatty acid esters and the like, and (vii) other suitable components such as glycerol triacetate and the like. Some of these, e.g. (iii) to (vii), display surfactant-like behaviour and may also be termed co-surfactants.

- (i) Suitable medium chain fatty acid triglycerides are neutral oils, e.g. neutral plant oils, in particular fractionated coconut oils, for example those known and commercially available under the trade names Captex, Myritol, Capmul, Captex, Neobee and Mazol; Miglyol 812 being the most preferred. Miglyol 812 is a fractionated coconut oil comprising caprylic-capric acid triglycerides and having a molecular weight = about 520 daltons. Fatty acid composition = C₆ max. about 3%, C₈ about 50 to 65%, C₁₀ about 30 to 45%, C₁₂ max 5%; acid no. = about 0.1; saponification no. about 330 to 345; iodine no. = max 1. Miglyol 812 is available from the Hüls company. These triglycerides are described in Fiedler, loc. cit., vol. , p. , incorporated herein by reference.
- (ii) Mixed mono-, di-, tri-glycerides preferably comprise mixtures of C₈ to C₁₀ or C₁₂₋₂₀ fatty acid mono-, di- and tri-glycerides, especially mixed C₁₆₋₁₈ fatty acid mono-, di- and triglycerides. The fatty acid component of the mixed mono-, di- and tri-glycerides may comprise both saturated and unsaturated fatty acid residues. Preferably however they are predominantly comprised of unsaturated fatty acid residues; in particular C₁₈ unsaturated fatty acid residues. Suitably the mixed mono-, di-, tri-glycerides comprise at least 60%, preferably at least 75%, more preferably at least 85% by weight of a C₁₈ unsaturated fatty acid (for example linolenic, linoleic and oleic acid) mono-, di- and tri-glycerides. Suitably the mixed mono-, di-, tri-glycerides comprise less than 20%, for example about 15% or 10% by weight or less, saturated fatty acid (for example palmitic and stearic acid) mono-, di- and tri-glycerides. Mixed mono-, di-, tri-glycerides are preferably predominantly comprised of mono- and di-glycerides; for example mono- and di-glycerides comprise at least 50%, more preferably at least 70% based on the total weight of the lipophilic phase or component. More preferably, the mono- and di-glycerides comprise at least 75% (for example about 80% or 85% by weight of the lipophilic component. Preferably monoglycerides comprise from about 25 to about 50%, based on the total weight of the lipophilic component, of the mixed mono-, di-, tri-glycerides. More preferably from about 30 to about 40% (for example 35 to 40%) monoglycerides are present. Preferably

diglycerides comprise from about 30 to about 60%, based on the total weight of the lipophilic component, of the mixed mono-, di-, tri-glycerides. More preferably from about 40 to about 55% (for example 48 to 50%) diglycerides are present. Triglycerides suitably comprise at least 5% but less than about 25 %, based on the total weight of the lipophilic component, of the mixed mono-, di-, tri-glycerides. More preferably from about 7.5 to about 15% (for example from about 9 to 12%) triglycerides are present. Mixed mono-, di-, tri-glycerides may be prepared by admixture of individual mono-, di- or tri-glycerides in appropriate relative proportion. Conveniently however they comprise transesterification products of vegetable oils, for example almond oil, ground nut oil, olive oil, peach oil, palm oil or, preferably, corn oil, sunflower oil or safflower oil and most preferably corn oil, with glycerol. Such transesterification products are generally obtained as described in GB 2 257 359 or WO 94/09211, the contents of which are incorporated herein by reference. Preferably some of the glycerol is first removed to give a "substantially glycerol free batch" when soft gelatine capsules are to be made. Purified transesterification products of corn oil and glycerol provide particularly suitable mixed mono-, di-, and tri-glycerides hereinafter referred to as "refined oil" and produced according to the description of GB 2 257 359 or WO 94/09211.

(iii) The lipophilic component may alternatively comprise e.g. a pharmaceutically acceptable oil, preferably with an unsaturated component such as a vegetable oil or fish oil. The lipophilic component may comprise suitable transesterified ethoxylated vegetable oils such as those obtained by reacting various natural vegetable oils (for example, maize oil, kernel oil, almond oil, ground nut oil, olive oil, soybean oil, sunflower oil, safflower oil and palm oil, or mixtures thereof) with polyethylene glycols that have an average molecular weight of from 200 to 800, in the presence of an appropriate catalyst. These procedures are known and an example is described in US Patent 3 288 824. Transesterified ethoxylated corn oil is particularly preferred. Transesterified ethoxylated vegetable oils are known and are commercially available under the trade name LABRAFIL (H. Fiedler, loc cit, vol 2, page 707, incorporated herein by reference). Examples are LABRAFIL M 2125 CS

(obtained from corn oil and having an acid number of less than about 2, a saponification number of 155 to 175, an HLB value of 3 to 4, and an iodine number of 90 to 110), and LABRAFIL M 1944 CS (obtained from kernel oil and having an acid number of about 2, a saponification number of 145 to 175 and an iodine number of 60 to 90). LABRAFIL M
5 2130 CS (which is a transesterification product of a C_{12-18} glyceride and polyethylene glycol and which has a melting point of about 35 to 40°C, an acid number of less than about 2, a saponification number of 185 to 200 and an iodine number of less than about 3) may also be used. The preferred transesterified ethoxylated vegetable oil is LABRAFIL M 2125 CS which can be obtained, for example, from Gattefossé, Saint-Priest Cedex, France.

10

(iv) Mono-, di- and mono/diglycerides, e.g. suitable C_8 to C_{10} fatty acid mono-, di- and tri-glycerides include Capmul MCM, Akoline MCM (from the Karlshamns company), Imwitor 308 and Imwitor 988 (from the Contensio company), which have an HLB value of about 3.8 (from the Contensio company), and especially esterification products of caprylic
15 or capric acid with glycerol. Preferred products of this class are e.g. those comprising or essentially consisting of caprylic/capric acid mono- and di-glycerides. C_8 to C_{10} mono-, di-glycerides having 6 to 10 mol-% polyoxyethylene groups, e.g. Softigen 767 (available from Contensio Chemicals). Monoglycerides, e.g. monooleate, glycerol monopalmitate and glycerol monostearate, for example as known and commercially available under the
20 trade names Myvatex, Myvaplex, and Myverol (Fiedler, loc. cit., 2, p. 1044, incorporated herein by reference) and acetylated, e.g. mono- and di-acetylated monoglycerides, for example as known under the trade name Myvacet (Fiedler, loc. cit., 2, p. 1043, incorporated herein by reference). Most preferred is Capmul MCM.

25 (v) Sorbitan fatty acid esters, e.g. of the type known and commercially available under the trade name Span, for example including sorbitan-monolaureyl ester (HLB 8.6), -monopalmityl ester (HLB 6.7), -monostearyl ester (HLB 4.7), -tristearyl ester (HLB 2.1), -monooleyl ester (HLB 4.3), and -trioleylesters (HLB 1.8) (Fiedler, loc. cit., 2, p. 1430, incorporated herein by reference).

(vi) Pentaerythriol fatty acid esters and polyalkylene glycol ethers and polyalkylene glycol ethers, for example pentaerythrite-dioleate, -distearate, -monolaurate, -polyglycol ether, and -monostearate as well as pentaerythrite-fatty acid esters (Fiedler, loc. cit., 2, p. 1158-
5 1160, incorporated herein by reference)

(vii) Other suitable components include glycerol triacetate or (1,2,3)-triacetin (Fiedler, loc. cit., 2, p. 1580, incorporated herein by reference); and sterols and derivatives thereof.

10 When present, the lipophilic component may comprise 5 to 85 % by weight of the composition of the invention, e.g. 10 to 85%; preferably 15 to 70 % by weight, more preferably about 20 to about 50 % by weight. Preferably the relative proportion of hydrophilic component(s), lipophilic component(s) and the surfactant(s) lie within the "microemulsion" region on a standard three way plot graph.

15

The present applicants also contemplate compositions of the invention in the form of microemulsion preconcentrate which may be free of refined fish oil and/or ethanol and/or transesterified ethoxylated vegetable oil.

20 The composition of the invention may further include additives or ingredients, for example antioxidants (such as ascorbyl palmitate, butyl hydroxy anisole (BHA), butyl hydroxy toluene (BHT) and tocopherols), antimicrobial agents, enzyme inhibitors, stabilizers, and/or preserving agents. These additives or ingredients may comprise about 0.05 to 5 %, preferably 0.05 to 1 % by weight of the total weight of the composition of the invention. The compositions of the
25 invention may also include sweetening or flavoring agents in an amount of up to about 2.5 or 5% by weight based on the total weight of the composition of the invention.

In another aspect the invention provides a process for the production of a composition of the invention, which process comprises bringing the active agent and (1) the hydrophilic

component, (2) the surfactant, and optionally (3) the lipophilic component into intimate admixture.

When required, the compositions of the invention are preferably compounded in unit dosage
5 form, for example by filling them into orally administrable capsule shells. The capsule shells
may be soft or hard gelatine capsule shells. Where the composition is in unit dosage form, each
unit dosage will suitably contain from 0.1 and 100 mg active agent, for example 0.1 mg, 1 mg,
5 mg, 10 mg, 15 mg, 25 mg or 50 mg, preferably between 10 and 100 mg of the active agent,
more preferably between 10 and 50 mg; for example 15, 20, 25, or 50 mg, more preferably
10 between 5 and 20 mg, most preferably 5 or 10 mg. Such unit dosage forms are suitable for
administration 1 to 5 times daily depending upon the particular purpose of therapy, the phase of
therapy and the like. However, if desired, the compositions may be in drink solution form and
may include water or any other aqueous system, e.g. fruit juice, milk, and the like, to provide
e.g. colloidal systems, suitable for drinking, e.g. with a dilution of from about 1:10 to about
15 1:100.

The compositions of the invention, e.g. those in the examples hereinafter, may show good
stability characteristics as indicated by standard stability trials, for example having a shelf life
stability of up to one, two or three years, and even longer. The compositions of this invention
20 in form of micellar precursors produce stable aqueous micelles, e.g. for up to one day or
longer. One group of compositions of the invention may be of high stability that are capable,
on addition to water, of providing aqueous microemulsions having an average particle size of
<2,000 Å (200 nm), e.g. <1,500 Å (150 nm).

25 The compositions of the invention exhibit especially advantageous properties when
administered orally; for example in terms of consistency and high level of bioavailability
obtained in standard bioavailability trials. These trials are performed in animals e.g. rats or
dogs or healthy volunteers using chromatographic methods, e.g. HPLC, or a specific or
nonspecific monoclonal kit to determine the level of the active agent in the blood. For

example, the composition of Example 5 administered p.o. to dogs may give surprisingly high C_{\max} values as detected by ELISA using a specific monoclonal antibody.

Pharmacokinetic parameters, for example absorption and blood levels, also become surprisingly
5 more predictable and problems in administration with erratic absorption may be eliminated or reduced. Additionally the compositions are effective with tenside materials, for example bile salts, being present in the gastro-intestinal tract. That is, the compositions may be fully dispersible in aqueous systems comprising such natural tensides and thus may be capable of providing microemulsion or aqueous micellar systems in situ which are stable and do not
10 exhibit precipitation of the active agent or other disruption of fine particulate structure. The function of the compositions upon oral administration may remain substantially independent of and/or unimpaired by the relative presence or absence of bile salts at any particular time or for any given individual. It also has been found that the compositions of this invention may reduce variability in inter- and intra-patient dose response.

15

The utility of the compositions of the invention may be observed in standard clinical tests in, for example, known indications of active agent at dosages giving therapeutically effective active agent blood levels. Any increased bioavailability of the compositions of the invention may be observed in standard animal tests and in clinical trials. For example, the utility of the
20 compositions of the invention in the treatment of social phobia may be established e.g. by administering these to subjects who have primary DSM-III-R diagnosis of social phobia, e.g. diagnosed by using the Structured Clinical Interview for DSM-III-R, e.g. in a 12 week open clinical trial. The treatment begins e.g. using 10 mg of an active agent daily and may be increased according to clinical response and side effects. Patients complete their self-report
25 measures at baseline and at e.g. weeks 4, 8, and 12. These measures include the fear of negative evaluation scale, social avoidance and distress scale, the social anxiety thoughts questionnaire, the fear questionnaire, the state-trait anxiety inventory, the Beck depression inventory, the social adjustment scale self-report, and the Sheehan disability scale. The responders e.g. at endpoint rate on the clinical global impression change by defining moderately

or markedly improved.

The dose of the active agent in the composition of the invention is of the same order as, or up to half, that used in known compositions containing the active agent. The compositions of the invention show activity at concentrations from about 0.1 mg to about 40 mg/day of active agent, preferably from about 0.1 mg to about 20 mg/day, e.g. most preferably from about 0.1 to about 1 mg/day of active agent for respiratory disease states and from about 5 to about 10 mg/day of active agent for CNS indications for a 75 kilogram mammal.

10 The compositions of the invention are particularly useful for treatment and prevention of the conditions disclosed in EP 0532456B1 and EP 0707006A2, the contents of which are incorporated herein by reference, which include treatment and prevention of central nervous system disorders, including depression, dysthymia, social phobia, panic disorder, and emesis. The compositions of the invention also are useful for the treatment of respiratory diseases, e.g. 15 asthma and chronic bronchitis. For these indications, the appropriate dosage will, of course, vary depending upon, for example, the particular composition of the invention employed, the host, the mode of administration, and the nature and severity of the conditions being treated.

A typical dose for Compound A is from 0.1 to 1 mg/day for asthma and chronic bronchitis and 20 from 5 to 10 mg/day of active agent for depression and social phobia for a 75 kilogram mammal, for the Example 5 form.

Thus in another aspect the present invention provides a method of treatment of a subject suffering from a disorder treatable with a piperidine substance P antagonist comprising 25 administering a therapeutically effective amount of a pharmaceutical composition of the invention to a subject in need of such treatment.

Examples

Following is a description by way of example only of compositions of this invention. Unless otherwise indicated, components are shown in % by weight based on each composition. Mean
5 particle sizes (diameters) are measured at 20° C using a Malvern Zetasizer.

All ingredients of the Examples are given in mg/capsule.

[illegible]

	Ex 1a ¹	Ex 1b ¹	Ex 2 ²	Ex 3 ²	Ex 4 ²	Ex 5 ²	Ex 6 ²	Ex 7 ²	Ex 8 ²
<u>3) Lipophilic component/ co-surfactant</u>									
Labrafil 2125				90.0	90.0				
Caprylic/capric acid glycerides (Capmul MCM)						170.0	218.5		
Propyleneglycol monocaprylate								133.0	136.0
Miglyol 812									
refined corn oil ³	185.0	185.0	90.0						
<u>4) Hydrophilic co- component</u>									
Ethanol abs	52.0	52.0	50.0	50.0	50.0	50.0	50.0	50.0	50.0
<u>Additive</u>									
DL-alpha tocopherol	0.5	0.5							
TOTAL	504.0	500.0	500.0	504.0	500.0	500.0	500.0	500.0	500.0
<u>mean particle size (nm)</u>	80-90	80-90	20-30	135-145	25-35	-	-	-	-

¹ Compound A is dissolved in (1) with stirring at room temperature and (2) and (3) are added to the obtained solution again with stirring. 0.5 ml portions of the obtained mixture are filled into size 1 hard gelatine capsules and sealed, e.g. using the Quali-Seal technique, or into soft gelatine capsules. In another embodiment of Examples 1a and 1b, Compound A is dispersed in a mixture of
5 components 1), 2) and 3), and combined with component 4).

² The carrier medium is prepared by mixing the components one with another. Compound A is then dissolved in the carrier medium by stirring.

10 ³ Refined oil = "refined glycerol-transesterified corn oil", substantially glycerol free, as described in GB 2 257 359 and WO 94/09211.

No phase separation or precipitation is observed for any of the above compositions 1 to 8 which are clear for 4 hours.

CLAIMS

1. A spontaneously dispersible pharmaceutical composition comprising a piperidine substance P antagonist.
- 5 2. A spontaneously dispersible pharmaceutical composition comprising a piperidine substance P antagonist and a carrier medium comprising
 - 1) a hydrophilic component and
 - 2) a surfactant.
- 10 3. A composition as claimed in claim 1 or 2 for oral administration.
4. A composition as claimed in any preceding claim comprising a lipophilic component.
- 15 5. A pharmaceutical composition comprising (2R,4S)-N-(1-(3,5-bis(trifluoromethyl)-benzoyl)-2-(4-chlorobenzyl)-4-piperidinyl)-quinoline-4-carboxamide as active agent and a carrier medium comprising
 - 1) a hydrophilic phase,
 - 2) a lipophilic phase, and
 - 20 3) a surfactant.
6. A pharmaceutical composition of any one of claims 1 to 4 comprising (2R,4S)-N-(1-(3,5-bis(trifluoromethyl)-benzoyl)-2-(4-chlorobenzyl)-4-piperidinyl)-quinoline-4-carboxamide.
- 25 7. A composition as claimed in any preceding claim in the form of a microemulsion concentrate.
8. A composition as claimed in any preceding claim wherein the hydrophilic component comprises triethyl citrate or propylene glycol, wherein the surfactant comprises a reaction

product of a natural or hydrogenated castor oil with ethylene oxide or a polyoxyethylene fatty acid ester, and wherein the lipophilic component comprises a transesterified ethoxylated vegetable oil, a C₈ to C₁₀ fatty acid mono-, di and/or tri-glyceride, a medium chain fatty acid triglyceride or a refined glycerol-transesterified corn oil.

5

9. A spontaneously dispersible pharmaceutical composition comprising about 0.05 to about 20 % by weight of (2R,4S)-N-(1-(3,5-bis(trifluoromethyl)-benzoyl)-2-(4-chlorobenzyl)-4-piperidinyl)-quinoline-4-carboxamide, about 5 to about 50 % by weight of a hydrophilic component, about 5 to about 80 % by weight of a surfactant, and optionally
10 about 5 to about 85 % by weight of a lipophilic component, all weights based on the total composition.
10. A composition as claimed in any preceding claim in the form of a microemulsion.
- 15 11. A composition according to any preceding claim in unit dosage form.
12. A composition according to claim 11 in soft or hard gelatin encapsulated form.
13. A method of treatment of a subject suffering from a disorder treatable with aa piperidine
20 substance P antagonist comprising administering a therapeutically effective amount of a pharmaceutical composition as claimed in any preceding claim to a subject in need of such treatment.

25

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 99/03623

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 A61K31/47 A61K9/107

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 0 707 006 A (CIBA GEIGY AG) 17 April 1996 (1996-04-17) cited in the application page 2, line 1-21 page 3, line 39-57 examples 3,9,10 claims 2-4,6	1,3,6,11
X	EP 0 826 684 A (HISAMITSU PHARMACEUTICAL CO) 4 March 1998 (1998-03-04) page 2, line 5-9 page 5, line 55,56 page 30-31; examples 4,6-11 claim 8	1,2



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

10 September 1999

Date of mailing of the international search report

17/09/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

La Gaetana, R

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 99/03623

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 0 722 722 A (OREAL) 24 July 1996 (1996-07-24) page 2, line 3,4 page 4, line 8-10 page 4, line 25-35 examples claims 1,5,12 -----	1,3
A	GB 2 278 780 A (SANDOZ LTD) 14 December 1994 (1994-12-14) page 3, line 8 - page 4, line 23 page 6, line 9 - page 9, line 18 page 12, line 3-24 page 16, line 11-13 claims -----	5-12

INTERNATIONAL SEARCH REPORT

International application No.

PCT/EP 99/03623

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Although claim 13 is directed to a method of treatment of the human body, the search has been carried out and based on the alleged effects of the composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 99/03623

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0707006 A	17-04-1996	AT 178602 T	15-04-1999
		AU 705851 B	03-06-1999
		AU 3317495 A	26-04-1996
		CA 2160444 A	15-04-1996
		CN 1133839 A	23-10-1996
		DE 59505575 D	12-05-1999
		FI 954829 A	15-04-1996
		HU 74103 A	28-11-1996
		JP 8176145 A	09-07-1996
		NO 954067 A	15-04-1996
		ZA 9508647 A	15-04-1996
EP 0826684 A	04-03-1998	AU 696948 B	24-09-1998
		AU 5120296 A	16-10-1996
		US 5886011 A	23-03-1999
		CA 2216701 A	03-10-1996
		WO 9630367 A	03-10-1996
EP 0722722 A	24-07-1996	FR 2728165 A	21-06-1996
		AT 171372 T	15-10-1998
		BR 9505381 A	18-11-1997
		CA 2165518 A	20-06-1996
		DE 69504949 D	29-10-1998
		DE 69504949 T	18-02-1999
		ES 2124983 T	16-02-1999
		HU 74948 A	28-03-1997
		JP 8208508 A	13-08-1996
		PL 311917 A	24-06-1996
		US 5744156 A	28-04-1998
GB 2278780 A	14-12-1994	GB 2315216 A, B	28-01-1998
		BE 1008329 A	02-04-1996
		CA 2124259 A	28-11-1994
		CH 686761 A	28-06-1996
		DE 4418115 A	01-12-1994
		ES 2098180 A	16-04-1997
		FR 2705566 A	02-12-1994
		IT 1272992 B	01-07-1997
		JP 7138161 A	30-05-1995
		US 5932243 A	03-08-1999

THIS PAGE BLANK (USPTO)